

Binary words containing infinitely many overlaps

James Currie

Department of Mathematics
University of Winnipeg
Winnipeg, Manitoba R3B 2E9 (Canada)
j.currie@uwinnipeg.ca

Narad Rampersad, Jeffrey Shallit

School of Computer Science
University of Waterloo
Waterloo, Ontario N2L 3G1 (Canada)
nrampersad@math.uwaterloo.ca
shallit@graceland.math.uwaterloo.ca

Submitted: Nov 16, 2005; Accepted: Sep 15, 2006; Published: Sep 22, 2006
Mathematics Subject Classifications: 68R15

Abstract

We characterize the squares occurring in infinite overlap-free binary words and construct various α power-free binary words containing infinitely many overlaps.

1 Introduction

If α is a rational number, a word w is an α power if there exists words x and x' , with x' a prefix of x , such that $w = x^n x'$ and $\alpha = n + |x'|/|x|$. We refer to $|x|$ as a *period* of w . An α^+ power is a word that is a β power for some $\beta > \alpha$. A word is α power-free (resp. α^+ power-free) if none of its subwords is an α power (resp. α^+ power). A 2 power is called a *square*; a 2^+ power is called an *overlap*.

Thue [18] constructed an infinite overlap-free binary word; however, Dekking [8] showed that any such infinite word must contain arbitrarily large squares. Shelton and Soni [17] characterized the overlap-free squares, but it is not hard to show that there are some overlap-free squares, such as 00110011, that cannot occur in an infinite overlap-free binary word. In this paper, we characterize those overlap-free squares that do occur in infinite overlap-free binary words.

Shur [16] considered the bi-infinite overlap-free and $7/3$ power-free binary words and showed that these classes of words were identical. There have been several subsequent papers [1, 10, 11, 14] that have shown various similarities between the classes of overlap-free binary words and $7/3$ power-free binary words. Here we contrast the two classes of words

by showing that there exist one-sided infinite $7/3$ power-free binary words containing infinitely many overlaps. More generally, we show that for any real number $\alpha > 2$ there exists a real number β arbitrarily close to α such that there exists an infinite β^+ power-free binary word containing infinitely many β powers.

All binary words considered in the sequel will be over the alphabet $\{0, 1\}$. We therefore use the notation \bar{w} to denote the *binary complement* of w ; that is, the word obtained from w by replacing 0 with 1 and 1 with 0.

2 Properties of the Thue-Morse morphism

In this section we present some useful properties of the *Thue-Morse morphism*; *i.e.*, the morphism μ defined by $\mu(0) = 01$ and $\mu(1) = 10$. It is well-known [12, 18] that the *Thue-Morse word*

$$\mathbf{t} = \mu^\omega(0) = 0110100110010110\dots$$

is overlap-free.

The following property of μ is easy to verify.

Lemma 1. *Let x and y be binary words. Then x is a prefix (resp. suffix) of y if and only if $\mu(x)$ is a prefix (resp. suffix) of $\mu(y)$.*

Brandenburg [6] proved the following useful theorem, which was independently rediscovered by Shur [16].

Theorem 2 (Brandenburg; Shur). *Let w be a binary word and let $\alpha > 2$ be a real number. Then w is α power-free if and only if $\mu(w)$ is α power-free.*

The following sharper version of one direction of this theorem (implicit in [10]) is also useful.

Theorem 3. *Suppose $\mu(w)$ contains a subword u of period p , with $|u|/p > 2$. Then w contains a subword v of length $\lceil |u|/2 \rceil$ and period $p/2$.*

Karhumäki and Shallit [10] gave the following generalization of the factorization theorem of Restivo and Salemi [15]. The extension to infinite words is clear.

Theorem 4 (Karhumäki and Shallit). *Let $x \in \{0, 1\}^*$ be α power-free, $2 < \alpha \leq 7/3$. Then there exist $u, v \in \{\epsilon, 0, 1, 00, 11\}$ and an α power-free $y \in \{0, 1\}^*$ such that $x = u\mu(y)v$.*

3 Overlap-free squares

Let

$$A = \{00, 11, 010010, 101101\}$$

and let

$$\mathcal{A} = \bigcup_{k \geq 0} \mu^k(A).$$

Pansiot [13] and Brlek [7] gave the following characterization of the squares in \mathbf{t} .

Theorem 5 (Pansiot; Brlek). *The set of squares in \mathbf{t} is exactly the set \mathcal{A} .*

We can use this result to prove the following.

Proposition 6. *For any position i , there is at most one square in \mathbf{t} beginning at position i .*

Proof. Suppose to the contrary that there exist distinct squares x and y that begin at position i . Without loss of generality, suppose that x and y begin with 0. Then by Theorem 5, $x = \mu^p(u)$ and $y = \mu^q(v)$, for some p, q and $u, v \in \{00, 010010\}$. Suppose $p \leq q$ and let $w = \mu^{q-p}(v)$. By Lemma 1, either u is a proper prefix of w or w is a proper prefix of u , neither of which is possible for any choice of $u, v \in \{00, 010010\}$. \square

The set \mathcal{A} does not contain all possible overlap-free squares. Shelton and Soni [17] characterized the overlap-free squares (the result is also attributed to Thue in [4]).

Theorem 7 (Shelton and Soni). *The overlap-free binary squares are the conjugates of the words in \mathcal{A} .*

Some overlap-free squares cannot occur in any infinite overlap-free binary word, as the following lemma shows.

Lemma 8. *Let $x = \mu^k(z)$ for some $k \geq 0$ and $z \in \{011011, 100100\}$. Then xa contains an overlap for all $a \in \{0, 1\}$.*

Proof. It is easy to see that $x = uvvuvv$ for some $u, v \in \{0, 1\}^*$, where u and v begin with different letters. Thus one of $uvvuvva$ or vva is an overlap. \square

We can characterize the squares that can occur in an infinite overlap-free binary word. Let

$$B = \{001001, 110110\}$$

and let

$$\mathcal{B} = \bigcup_{k \geq 0} \mu^k(B).$$

Theorem 9. *The set of squares that can occur in an infinite overlap-free binary word is $\mathcal{A} \cup \mathcal{B}$. Furthermore, if \mathbf{w} is an infinite overlap-free binary word containing a subword $x \in \mathcal{B}$, then \mathbf{w} begins with x and there are no other occurrences of x in \mathbf{w} .*

Proof. Let \mathbf{w} be an infinite overlap-free binary word beginning with a square $yy \notin \mathcal{A} \cup \mathcal{B}$. Suppose further that yy is a smallest such square that can be extended to an infinite overlap-free word. If $|y| \leq 3$, then $yy \notin \mathcal{A} \cup \mathcal{B}$ is one of 011011 or 100100, neither of which can be extended to an infinite overlap-free word by Lemma 8.

We assume then that $|y| > 3$. Since, by Theorem 7, yy is a conjugate of a word in \mathcal{A} , we have two cases.

Case 1: $yy = \mu(zz)$ for some $z \in \{0, 1\}^*$. By Theorem 4, $\mathbf{w} = \mu(zz\mathbf{w}')$ for some infinite \mathbf{w}' , where $zz\mathbf{w}'$ is overlap-free. Thus zz is a smaller square not in $\mathcal{A} \cup \mathcal{B}$ that can be extended to an infinite overlap-free word, contrary to our assumption.

Case 2: $yy = a\mu(zz')\bar{a}$ for some $a \in \{0, 1\}$ and $z, z' \in \{0, 1\}^*$. By Theorem 4, yy is followed by a in \mathbf{w} , and so yya is an overlap, contrary to our assumption.

Since both cases lead to a contradiction, our assumption that $yy \notin \mathcal{A} \cup \mathcal{B}$ must be false.

To see that each word in $\mathcal{A} \cup \mathcal{B}$ does occur in some infinite overlap-free binary word, note that Allouche, Currie, and Shallit [2] have shown that the word $\mathbf{s} = 001001\bar{\mathbf{t}}$ is overlap-free. Now consider the words $\mu^k(\mathbf{s})$ and $\mu^k(\bar{\mathbf{s}})$, which are overlap-free for all $k \geq 0$.

Finally, to see that any occurrence of $x \in \mathcal{B}$ in \mathbf{w} must occur at the beginning of \mathbf{w} , we note that by an argument similar to that used in Lemma 8, ax contains an overlap for all $a \in \{0, 1\}$, and so x occurs at the beginning of \mathbf{w} . \square

4 Words containing infinitely many overlaps

In this section we construct various infinite α power-free binary words containing infinitely many overlaps. We begin by considering the infinite $7/3$ power-free binary words.

Proposition 10. *For all $p \geq 1$, an infinite $7/3$ power-free word contains only finitely many occurrences of overlaps with period p .*

Proof. Let \mathbf{x} be an infinite $7/3$ power-free word containing infinitely many overlaps with period p . Let $k \geq 0$ be the smallest integer satisfying $p \leq 3 \cdot 2^k$. Suppose \mathbf{x} contains an overlap w with period p starting in a position $\geq 2^{k+1}$. Then by Theorem 4, we can write

$$\mathbf{x} = u_1\mu(u_2) \cdots \mu^{k-1}(u_k)\mu^k(\mathbf{y}),$$

where each $u_i \in \{\epsilon, 0, 1, 00, 11\}$. The overlap w occurs as a subword of $\mu^k(\mathbf{y})$. By Lemma 3, \mathbf{y} contains an overlap with period $p/2^k \leq 3$. But any overlap with period ≤ 3 contains a $7/3$ power. Thus, \mathbf{x} contains a $7/3$ power, a contradiction. \square

The following theorem provides a striking contrast to Shur's result [16] that the bi-infinite $7/3$ power-free words are overlap-free.

Theorem 11. *There exists a $7/3$ power-free binary word containing infinitely many overlaps.*

Proof. We define the following sequence of words: $A_0 = 00$ and $A_{n+1} = 0\mu^2(A_n)$, $n \geq 0$. The first few terms in this sequence are

$$\begin{aligned} A_0 &= 00 \\ A_1 &= 001100110 \\ A_2 &= 001100110100110011001011001101001100110010110 \\ &\vdots \end{aligned}$$

We first show that in the limit as $n \rightarrow \infty$, this sequence converges to an infinite word \mathbf{a} . It suffices to show that for all n , A_n is a prefix of A_{n+1} . We proceed by induction on n . Certainly, $A_0 = 00$ is a prefix of $A_1 = 0\mu^2(00) = 001100110$. Now $A_n = 0\mu^2(A_{n-1})$, $A_{n+1} = 0\mu^2(A_n)$, and by induction, A_{n-1} is a prefix of A_n . Applying Lemma 1, we see that A_n is a prefix of A_{n+1} , as required.

Note that for all n , A_{n+1} contains $\mu^{2n}(A_1)$ as a subword. Since A_1 is an overlap with period 4, $\mu^{2n}(A_1)$ contains 2^{2n} overlaps with period 2^{2n+2} . Thus, \mathbf{a} contains infinitely many overlaps.

We must show that \mathbf{a} does not contain a $7/3$ power. It suffices to show that A_n does not contain a $7/3$ power for all $n \geq 0$. Again, we proceed by induction on n . Clearly, $A_0 = 00$ does not contain a $7/3$ power. Consider $A_{n+1} = 0\mu^2(A_n)$. By induction, A_n is $7/3$ power-free, and by Theorem 2, so is $\mu^2(A_n)$. Thus, if A_{n+1} contains a $7/3$ power, such a $7/3$ power must occur as a prefix of A_{n+1} . Note that A_{n+1} begins with 00110011. The word 00110011 cannot occur anywhere else in A_{n+1} , as that would imply that A_{n+1} contained a cube 000 or 111, or the $5/2$ power 1001100110. If A_{n+1} were to begin with a $7/3$ power with period ≥ 8 , it would contain two occurrences of 00110011, contradicting our earlier observation. We conclude that the period of any such $7/3$ power is less than 8. Checking that no such $7/3$ power exists is now a finite check and is left to the reader. \square

In fact, we can prove the following stronger statement.

Theorem 12. *There exist uncountably many $7/3$ power-free binary words containing infinitely many overlaps.*

Proof. For a finite binary sequence b , we define an operator g_b on binary words recursively by

$$\begin{aligned} g_\epsilon(w) &= w \\ g_{0b}(w) &= \mu^2(g_b(w)) \\ g_{1b}(w) &= 0\mu^2(g_b(w)). \end{aligned}$$

Note that $g_b(0)$ always starts with a 0, so that for any finite binary words p and b , $g_p(0)$ is always a prefix of $g_{pb}(0)$. Since $g_0(0)$ is not a prefix of $g_1(0)$, $g_{p0}(0)$ is not a prefix of $g_{p1}(0)$ for any p , so that distinct b give distinct words. Given an infinite binary sequence $\mathbf{b} = b_1b_2b_3 \cdots$ where the $b_i \in \{0, 1\}$, define an infinite binary sequence $w_{\mathbf{b}}$ to be the limit of

$$g_\epsilon(00), g_{b_1}(00), g_{b_1b_2}(00), g_{b_1b_2b_3}(00), \dots$$

By an earlier argument, each $w_{\mathbf{b}}$ is $7/3$ power-free. Since $g_1(00) = 001100110$ is an overlap, $g_{b_1}(00) = g_b(001100110)$ ends with an overlap for any finite word b . Thus, each 1 in \mathbf{b} introduces an overlap in $w_{\mathbf{b}}$. Since uncountably many binary sequences contain infinitely many 1's, uncountably many of the $w_{\mathbf{b}}$ are $7/3$ power-free words containing infinitely many overlaps. \square

Next, we show that the sequence \mathbf{a} constructed in the proof of Theorem 11 is an automatic sequence (in the sense of [3]).

Proposition 13. *The sequence \mathbf{a} is 4-automatic.*

Proof. We show that $\mathbf{a} = g(h^\omega(0))$, where h and g are the morphisms defined by

$$\begin{array}{ll} h(0) = 0134 & g(0) = 0 \\ h(1) = 2134 & g(1) = 0 \\ h(2) = 3234 & g(2) = 0 \\ h(3) = 2321 & g(3) = 1 \\ h(4) = 3421 & g(4) = 1. \end{array} \quad \text{and}$$

We make some observations concerning 2-letter subwords: The sequence $h^\omega(0)$ clearly does not contain any of the words 11, 14, 22, 24, 31, 33, 41 or 44. In fact, neither 12 nor 43 appears as a subword either: Words 12 and 43 do not appear internally in $h(i)$, $0 \leq i \leq 4$; therefore, if 43 appears in $h^n(0)$, it must 'cross the boundary' in one of $h(12)$, $h(14)$, $h(22)$ or $h(24)$. Since 14, 22 and 24 do not appear in $h^\omega(0)$, word 43 can only appear in $h^n(0)$ as a descendant of a subword 12 in $h^{n-1}(0)$. However, the situation is symmetrical; word 12 can only appear in $h^n(0)$ as a descendant of a subword 43 in $h^{n-1}(0)$. By induction, neither 43 nor 12 ever appears.

The point of the previous paragraph is that

$$\begin{array}{l} h(0) \text{ always occurs in the context } h(0)2 \\ h(1) \text{ always occurs in the context } h(1)2 \\ h(2) \text{ always occurs in the context } h(2)2 \\ h(3) \text{ always occurs in the context } h(3)3 \\ h(4) \text{ always occurs in the context } h(4)3 \end{array}$$

The word $h^\omega(0)$ can thus be parsed in terms of a new morphism f :

$$\begin{array}{ll} f(0) = 1342 \\ f(1) = 1342 \\ f(2) = 2342 \\ f(3) = 3213 \\ f(4) = 4213. \end{array}$$

The parsing in terms of f works as follows: If we write $h^\omega(0) = 0w$, then $w = f(0w)$. It is useful to rewrite this relation in terms of the finite words $h^n(0)$. For non-negative integer n let x_n be the unique letter such that $h^n(0)x_n$ is a prefix of $h^\omega(0)$. Thus $x_0 = 1$, $x_1 = 2$, etc. We then have

$$h^n(0)x_n = 0f(h^{n-1}(0)), \quad n \geq 1. \quad (1)$$

Since for all $a \in \{0, 1, 2, 3, 4\}$, $g(f(a)) = \mu^2(g(a))$, we have $g(f(u)) = \mu^2(g(u))$ for all words u . Therefore, applying g to (1)

$$\begin{aligned} g(h^n(0)x_n) &= g(0f(h^{n-1}(0))) \\ &= g(0)g(f(h^{n-1}(0))) \\ &= 0\mu^2(g(h^{n-1}(0))), \quad n \geq 1. \end{aligned}$$

From this relation we show by induction that A_n is the prefix of $g(h^{n+1}(0))$ of length $(4^{n+1} + 3 \cdot 4^n - 1)/3$. Certainly, $A_0 = 00$ is the prefix of length 2 of $g(h(0)) = 0011$. Consider $A_n = 0\mu^2(A_{n-1})$. We can assume inductively that A_{n-1} is the prefix of $g(h^n(0))$ of length $(4^n + 3 \cdot 4^{n-1} - 1)/3$. Writing $g(h^n(0)) = A_{n-1}z$ for some z , we have

$$\begin{aligned} g(h^{n+1}(0)x_{n+1}) &= 0\mu^2(g(h^n(0))) \\ &= 0\mu^2(A_{n-1}z) \\ &= A_n\mu^2(z), \end{aligned}$$

for some x_{n+1} , whence A_n is a prefix of $g(h^{n+1}(0))$. Since $|A_n| = 4|A_{n-1}| + 1$, we have $|A_n| = (4^{n+1} + 3 \cdot 4^n - 1)/3$, as required. \square

The result of Theorem 11 can be strengthened even further.

Theorem 14. *For every real number $\alpha > 2$ there exists a real number β arbitrarily close to α , such that there is an infinite β^+ power-free binary word containing infinitely many β powers.*

Proof. Let $s \geq 3$ be a positive integer, and let $r = \lfloor \alpha + 1 \rfloor$. Let t be the largest positive integer such that $r - t/2^s > \alpha$, and such that the word obtained by removing a prefix of length t from $\mu^s(0)$ begins with 00. Let $\beta = r - t/2^s$. Since $\alpha \geq r - 1$, we have $t < 2^s$. Also, $\mu^3(0) = 01101001$ and $\mu^3(1) = 10010110$ are of length 8, and both contain 00 as a subword; it follows that $|\alpha - \beta| \leq 8/2^s$, so that by choosing large enough s , β can be made arbitrarily close to α .

We construct sequences of words A_n , B_n and C_n . Define $C_0 = 00$. For each $n \geq 0$:

1. Let $A_n = 0^{r-2}C_n$.
2. Let $B_n = \mu^s(A_n)$.
3. Remove the first t letters from B_n to obtain a new word C_{n+1} beginning with 00.

Since each A_n begins with the r power 0^r , each $B_n = \mu^s(A_n)$ begins with an r power of period 2^s . Removing the first t letters ensures that C_{n+1} commences with an $(r2^s - t)/2^s$ power, viz., a β power. The limit of the C_n gives the desired infinite word. Let us check that this limit exists:

Let w be the word consisting of the first t letters of $\mu^s(0)$. Since all the A_n commence with 0 by construction, all the B_n commence with $\mu^s(0)$, and hence with w . This means that $B_n = wC_{n+1}$ for each n .

We show that A_n is always a prefix of A_{n+1} by induction. Certainly A_0 is a prefix of A_1 . Assume that A_{n-1} is a prefix of A_n . Since $A_n = 0^{r-2}C_n$ and $A_{n+1} = 0^{r-2}C_{n+1}$, A_n is a prefix of A_{n+1} if C_n is a prefix of C_{n+1} . Since $B_{n-1} = wC_n$ and $B_n = wC_{n+1}$, C_n is a prefix of C_{n+1} if B_{n-1} is a prefix of B_n . By Lemma 1, B_{n-1} is a prefix of B_n if A_{n-1} is a prefix of A_n , which is our inductive assumption. We conclude that A_n is a prefix of A_{n+1} .

It follows that C_n is a prefix of C_{n+1} for $n \geq 0$, so that the limit of the C_n exists. It will thus suffice to prove the following claim:

Claim: The A_n , B_n and C_n satisfy the following:

1. The word C_n contains no β^+ powers.
2. The only β^+ power in A_n is 0^r .
3. Any β^+ powers in B_n appear only in the prefix $\mu^s(0^r)$.

Certainly C_0 contains no β^+ powers, and since $\beta > r - 1$, the only β^+ power in A_0 is 0^r . Suppose then that the claim holds for A_n and C_n .

Now suppose that $B_n = \mu^s(0^{r-2})\mu^s(C_n)$ contains a β^+ power u with period p . Since C_n contains no β^+ powers, Theorem 2 ensures that $\mu^s(C_n)$ contains no β^+ powers. We can therefore write $B_n = xuy$ where $|x| < |\mu^s(0^{r-2})|$. In other words, u overlaps $\mu^s(0^{r-2})$ from the right. By Theorem 3, the preimage of B_n under μ , i.e., $\mu^{s-1}(A_n)$, contains a β^+ power of length at least $|u|/2$ and period $p/2$. In fact, iterating this argument, A_n contains a β^+ power of period $p/2^s$ of length at least $|u|/2^s$. Since the only β^+ power in A_n is 0^r , with period 1, we see that $p/2^s = 1$, whence $p = 2^s$ and $|u| \leq r2^s$.

Recall that B_n has a prefix $\mu^s(0^r)$ which also has period 2^s , and that this prefix is overlapped by u . It follows that all of xu is a β^+ power with period $p = 2^s$. However, as just argued, this means that $|xu| \leq r2^s = |\mu^s(0^r)|$, so that u is contained in $\mu^s(0^r)$ and part 3 of our claim holds for B_n . We now show that parts 1 and 2 hold for C_{n+1} and A_{n+1} respectively, and the truth of our claim will follow by induction.

Part 1 follows immediately from part 3.

Now suppose that A_{n+1} contains a β^+ power u . Recall that $A_{n+1} = 0^{r-2}C_{n+1}$, and C_{n+1} begins with 00 , but contains no β^+ powers. It follows that u is not a subword of C_{n+1} . Therefore, 000 must be a prefix of u . If $u = 0^q$ for some integer q , then $q \leq r$ by the construction of A_{n+1} , and

$$r \geq q > \beta > \alpha > r - 1.$$

This implies that $q = r$, and $u = 0^r$, as claimed. If we cannot write $u = 0^q$, then $|u|_1 \geq 1$. Because u is a 2^+ power, 000 must appear twice in u with a 1 lying somewhere between the

two appearances. This implies that 000 is a subword of C_{n+1} , and hence of $B_n = \mu^s(A_n)$. However, no word of the form $\mu(w)$ contains 000. This is a contradiction. \square

We conclude by presenting the following open problem.

Does there exist a characterization (in the sense of [5, 9]) of the infinite $7/3$ power-free binary words?

5 Acknowledgments

Thanks to the referee for pointing out Brandenburg's proof of Theorem 2.

References

- [1] A. Aberkane, J. Currie, "Attainable lengths for circular binary words avoiding k powers", *Bull. Belg. Math. Soc. Simon Stevin*, 2004, to appear.
- [2] J.-P. Allouche, J. Currie, J. Shallit, "Extremal infinite overlap-free words", *Electron. J. Combin.* **5** (1998), #R27.
- [3] J.-P. Allouche, J. Shallit, *Automatic Sequences: Theory, Applications, Generalizations*, Cambridge, 2003.
- [4] J. Berstel, "Axel Thue's work on repetitions in words". In P. Leroux, C. Reutenauer, eds., *Séries formelles et combinatoire algébrique*, Publications du LaCIM, pp 65–80, UQAM, 1992.
- [5] J. Berstel, "A rewriting of Fife's theorem about overlap-free words". In J. Karhumäki, H. Maurer, G. Rozenberg, eds., *Results and Trends in Theoretical Computer Science*, Vol. 812 of *Lecture Notes in Computer Science*, pp. 19–29, Springer-Verlag, 1994.
- [6] F.-J. Brandenburg, "Uniformly growing k -th power-free homomorphisms", *Theoret. Comput. Sci.* **23** (1983), 69–82.
- [7] S. Brlek, "Enumeration of factors in the Thue-Morse word", *Discrete Appl. Math.* **24** (1989), 83–96.
- [8] F. M. Dekking, "On repetitions in binary sequences", *J. Comb. Theory Ser. A* **20** (1976), 292–299.
- [9] E. Fife, "Binary sequences which contain no BBb ", *Trans. Amer. Math. Soc.* **261** (1980), 115–136.
- [10] J. Karhumäki, J. Shallit, "Polynomial versus exponential growth in repetition-free binary words", *J. Combin. Theory Ser. A* **104** (2004), 335–347.

- [11] R. Kolpakov, G. Kucherov, Y. Tarannikov, “On repetition-free binary words of minimal density”, WORDS (Rouen, 1997), *Theoret. Comput. Sci.* **218** (1999), 161–175.
- [12] M. Morse, G. Hedlund, “Unending chess, symbolic dynamics, and a problem in semi-groups”, *Duke Math. J.* **11** (1944), 1–7.
- [13] J. J. Pansiot, “The Morse sequence and iterated morphisms”, *Inform. Process. Lett.* **12** (1981), 68–70.
- [14] N. Rampersad, “Words avoiding $\frac{7}{3}$ -powers and the Thue-Morse morphism”, *Internat. J. Found. Comput. Sci.* **16** (2005), 755–766.
- [15] A. Restivo, S. Salemi, “Overlap free words on two symbols”. In M. Nivat, D. Perrin, eds., *Automata on Infinite Words*, Vol. 192 of *Lecture Notes in Computer Science*, pp. 198–206, Springer-Verlag, 1984.
- [16] A. M. Shur, “The structure of the set of cube-free \mathbb{Z} -words in a two-letter alphabet” (Russian), *Izv. Ross. Akad. Nauk Ser. Mat.* **64** (2000), 201–224. English translation in *Izv. Math.* **64** (2000), 847–871.
- [17] R. Shelton, R. Soni, “Chains and fixing blocks in irreducible binary sequences”, *Discrete Math.* **54** (1985), 93–99.
- [18] A. Thue, “Über die gegenseitige Lage gleicher Teile gewisser Zeichenreihen”, *Kra. Vidensk. Selsk. Skrifter. I. Math. Nat. Kl.* **1** (1912), 1–67.